

Q: Find the MLE of $\theta = (\mu, \sigma^2)$. Approximate the distribution of $\hat{\theta}$

The Y_1, Y_2, \dots, Y_n are i.i.d. $N(\mu, \sigma^2)$,

Let $T_1 = \Delta\mu$ and $T_2 = \Delta\sigma^2$, $T = (T_1, T_2)$

We already know that \hat{T}_{MLE} is approximately normal with covariance

$$I^{-1}(T_0)/n = \begin{bmatrix} \frac{T_2}{n} & 0 \\ 0 & \frac{2T_2^2}{n} \end{bmatrix} \quad \text{where } T_2 = \sigma^2$$

To get the covariance for MLE of (μ, σ^2) , we can use the Δ -method

$$\mu = g_1(T) = \mu + \frac{\sigma^2}{2} = \frac{T_1}{\Delta} + \frac{T_2}{2\Delta}$$

$$\sigma^2 = g_2(T) = \frac{T_2}{\Delta}$$

By invariance prop.

$$\hat{\mu}_{MLE} = g_1(\hat{T}_{MLE}) = \frac{\hat{T}_1}{\Delta} + \frac{\hat{T}_2}{2\Delta}$$

$$\hat{\sigma}_{MLE}^2 = \frac{\hat{T}_2}{\Delta}$$

The matrix G in this case: $m=2$ $p=2$, so $G_{2 \times 2}$

$$G = \begin{bmatrix} \frac{\partial g_1(T)}{\partial T_1} & \frac{\partial g_1(T)}{\partial T_2} \\ \frac{\partial g_2(T)}{\partial T_1} & \frac{\partial g_2(T)}{\partial T_2} \end{bmatrix} = \begin{bmatrix} \frac{1}{\Delta} & \frac{1}{2\Delta} \\ 0 & \frac{1}{\Delta} \end{bmatrix}$$

thus $GI^{-1}(T)G^T/n$

$$= \begin{bmatrix} \frac{T_2}{n\Delta^2} + \frac{T_2^2}{2n\Delta^2} & \frac{T_2^2}{n\Delta^2} \\ \frac{T_2^2}{n\Delta^2} & \frac{2T_2^2}{n\Delta^2} \end{bmatrix} \quad \text{note } T = n\Delta$$

$$\sigma^2 = \frac{T_2}{\Delta}$$

$$= \begin{bmatrix} \frac{\sigma^2}{T} + \frac{\sigma^4}{2n} & \frac{\sigma^4}{n} \\ \frac{\sigma^4}{n} & \frac{2\sigma^4}{n} \end{bmatrix}$$

The MLE for (α, σ^2) is approximately bivariate normal distribution based on this covariance matrix

Q: What does this result suggest regarding how large n should be chosen for the best estimation of σ^2 ?

What about for α ?

$V(\hat{\sigma}^2) = \frac{2\sigma^4}{n} \Rightarrow$ in order to estimate σ^2 more precisely, we can sample $[0, T]$ more finitely.

But for α , its estimation is limited by T :

$$V(\hat{\alpha}) = \frac{\sigma^2}{T} + \frac{\sigma^4}{2n} \approx \frac{\sigma^2}{T} \quad \text{for large } n, \text{ holding } T \text{ constant}$$

so we need a larger time interval to estimate α better.

Typical value of σ is around 0.2, so if we want $SE(\hat{\alpha}) \leq 0.01$

we need $T \geq 400$ years